

Subdeltoid Bursa Tuberculosis with Rice Bodies Formation: Case Report and Review of Literature

Milind.V .Pimprikar¹, Aashay L. Kekatpure²

What to Learn from this Article?

Presentation and Management of Tuberculous Subdeltoid Bursitis.

Abstract

Introduction: We describe a rare case of a patient with unilateral musculoskeletal manifestation of tuberculosis presented as bursitis of the left shoulder with rice bodies, without coexisting active tuberculosis or tuberculosis in the previous history.

Case Report: A 21 year old patient was examined, who complained of pain and swelling in the left shoulder for 2 years. MRI showed a large amount of rice bodies with joint effusion in the left shoulder with intact rotator cuff. The histological examination showed a tuberculosis-specific inflammatory response with giant cells and epithelioid granulomas. Arthroscopic debridement and removal of the loose bodies was done. A brief summary of the literature is given.

Conclusion: We report a unique case of tuberculous subdeltoid bursitis with rice bodies formation in absence of any other concomitant focus of tuberculous infection, managed with arthroscopic debridement and anti-tuberculous regimen with a long follow up of twelve months.

Keywords: Subdeltoid Bursitis; Rice bodies; Arthroscopy

Introduction

Rice body formation is commonly observed in the joint and tendon sheaths among patients with rheumatoid arthritis [1,2], however only a few cases with rice bodies in subdeltoid bursa of tubercular origin have been mentioned in the literature [3,4]. There are very few reports [5] about the arthroscopic management of cases with rice bodies in subdeltoid bursa with a long term follow up. The authors report the rare case of a patient with musculoskeletal manifestation of tuberculosis presented as bursitis of the left shoulder

with rice bodies without coexisting active tuberculosis or tuberculosis in the previous history managed with arthroscopic debridement with follow up of one year.

Case Report

A 21 years young boy with 2 years history of diffuse swelling of left shoulder was examined (Non dominant side). He was an athlete. Pain was gradual in onset. The pain was not aggravated by activities of daily living but terminal rotations were painful. There was no history of constitutional symptoms. There was no history of tuberculosis or any

Author's Photo Gallery



Dr. Milind.V .Pimprikar



Dr. Aashay L. Kekatpure

Reviewer's Photo Gallery



Dr. Ashwin Kasturi



Dr. Santosh Banshelkikar

¹Director And Senior Consultant ADTOOS Clinics & Pimprikar Hospital Nasik. India.

²Consultant Shoulder And Elbow Surgeon ADTOOS Clinics & Pimprikar Hospital Nasik. India.

Address of Correspondence

Dr. Aashay .L.Kekatpure, Pimprikar Hospital, Govind Nagar, Chowk No.5, Behind Prakash Hotel, Mumbai Naka, Nasik-400009. India.
E-mail: dr.aashayk@gmail.com




Figure 1: T2 Fat Suppressed Coronal Image Showing Intermediate Intensity Multiple Rice Bodies.



Figure 2: T2 Fat Suppressed Axial Image Showing Intermediate Intensity Multiple Rice Bodies.



Figure 3: Arthroscopic Image of Subdeltoid Rice Bodies.

other major illness in the past.

He was a febrile with vital parameter within normal limits. There were no signs of acute or chronic inflammation. There was diffuse swelling over the shoulder. No point tenderness was elicited; external and internal rotations of the shoulder were terminally restricted.

The WBC count was 7,900/cmm. With lymphocytes being 36%, His ESR (erythrocyte Sedimentation Rate) was raised (61mm). C-reactive protein study was positive. Rheumatoid factor and HIV studies were negative.

Radiograph showed no abnormality of the humeral head. Chest x-ray did not show any evidence of healed primary lesion. Magnetic resonance imaging (MRI) scans [Fig 1 & 2] showed moderate joint effusion with multiple loose bodies on T2 weighted image & signal changes of humeral head near the synovial reflection along posterior aspect. The T1 weighted images showed homogenous images. The patient underwent arthroscopic debridement [Fig 3] for removal of loose bodies some of which were attached to the synovium. The loose bodies resembled rice bodies ranging from 3 to 10 mm length [Fig 4].

Histo-pathological analysis confirmed caseous necrosis and ziehl neelsen staining of the fluid from the bursa isolated mycobacterium tuberculosis. Microscopically they consisted of compact fibrous material; however the articular cartilage did not show evidence of tuberculosis.

The patient was treated with anti-tuberculosis medication for 6 months and he is disease free for the last 12 months.

Discussion

The formation of intra-articular rice bodies was first described in tuberculous arthritis. These nodules are a common finding in rheumatoid bursitis and arthritis; they are rare in other arthropathies[6]. The pathogenesis of these rice bodies remains obscure. The main theories depict the origin of the rice bodies from the synovial fluid due to aggregation of the fibronectin /fibrin[7]. A different theory proposes them to be of the synovial origin. The underlying

disease condition leads to micro infraction of the synovium which progresses to sloughing, and then the fibrin covering the infraction tissue as described by Cheung [8].

The MRI findings in our patient revealed the rice bodies as intermediate intensity images on the T1 and T2 weighted images. They were better delineated on the T2 weighted images than T1 weighted images, on which they appeared homogenous. On arthroscopy they seemed attached to the synovial lining supporting the description of their origin by Cheung. Macroscopically they appeared similar to the rice bodies arising in rheumatoid arthritis. On microscopic analysis they were composed of fibrous tissue.

Tuberculous bursitis is always described secondary to some other primary focus of infection in the bone or the nearby joint [9]. Tuberculous involvement have been described in the superficially situated bursa such as the olecranon and the pre-patellar bursa[10-12] . History of trauma and direct transmission has been thought as the underlying cause. Hematogenous spread is proposed as the cause of deep seated bursa involvement[13]. For the hematogenous spread to occur there should be a primary focus of infection or a healed primary lesion in the lungs. In our patient there was no history of trauma and the chest x ray did not show any focus of infection .

So far very few studies have been published on sub-deltoid bursitis and its arthroscopic management with a long term follow up. Jaovisidha et al [4] has published a case series of 3 cases with subdeltoid bursitis. Alkalay et al [3] has reported a case of patient with 30 year history of tuberculous subdeltoid bursitis. Kim et al [5] reported a case with subdeltoid bursitis in a 41 year old woman. Arthroscopic debridement and biopsy of the subdeltoid bursa were performed along with biopsy of the tissue. Patient underwent anti tuberculosis treatment for 6 months , she was symptom free for 18 months.

The occurrence of subdeltoid bursa tuberculosis without any preexisting history is rare. Its should be ruled out as a cause of insidious shoulder pain. The results with arthroscopic



Figure 4: Rice Bodies Gross Anatomical Presentation.

debridement of the subdeltoid tuberculosis shows good results. More cases and longer follow up can help in determining the outcome. To date no recurrence has been reported after the arthroscopic debridement and antituberculosis medication.[5,13].

Conclusion

We report a unique case of tuberculous subdeltoid bursitis with rice body formation in absence of any other concomitant focus of tuberculous infection, managed with arthroscopic debridement and category III anti-tuberculous regimen according to DOTS with a long follow up of twelve months.

Clinical Message

Possibility of tuberculosis of subdeltoid bursa in absence of a primary focus should be ruled out.

References

1. Hiroyuki Nagasawa, et al Tenosynovitis with rice body formation in a non-tuberculosis patient: A case report *Ups J Med Sci.* 2009
2. Karthikeyan Iyengar ,et al Bilateral recurrent wrist flexor tenosynovitis and rice body formation in a patient with sero-negative rheumatoid arthritis: A case report and review of literature *Int J Surg Case Rep.* 2011; 2(7): 208–211.
3. Alkalay I, Kaufman T, Suprum H. Tuberculosis of the subdeltoid bursa. *Acasereport Isr j Med Sci* 1980;16:853-5
4. Jaovisidha S, Chen C, Ryu KN, Siriwong-pairat P, Pekan P, Sartoris DJ, et. al Tuberculous tenosynovitis and bursitis: imaging findings in 21 cases. *Radiology* 1996;201:507-13
5. Kim et al Tuberculous subdeltoid Bursitis with rice bodies. *Yonsei Medical Journal* 2002;43:539-42
6. Poper J. Rice – Bodies, Synovial Debris, and joint Lavage. *BR. J Rheumatol* 1985;24:1-2
7. Poper AJ, Scott DL, Wainwright AC, Walton KW, Williamson N, Chapman JH. Frequency of occurrence, mode of develop. And significance of rice bodies in rheumatoid jt. *Ann Rheum Dis* 1982;41:109-17
8. Cheung HS, Ryan LM, Kozin F, McCarty DJ, Synovial origins of rice bodies in joint, *fluid Arthritis Rheum* 1980;23:72-6
9. Watts HG, Lifeso RM. Tuberculosis of bone and joints. *J Bone Joint Surg Am* 1996;78:288-98
10. Dawson DJ, Blacklock ZM, Ashdown LR, Bittger EC. *Mycobacterium Asiaticum* as the probable causative agent in a case of olecranon Bursitis. *J. Clin Microbiol* 1995;33:1042-3
11. Holder SF, Hopson CN, Vonkuster LC. tuberculous arthritis of the elbow presenting as chronic bursitis of the olecranon . A case report. *J Bone Joint Surg AM* 1985;67:1127-9
12. Schickendantz MS, Watson JT, Mycobacterial prepatellar bursitis. *Clin Orthop* 1990;258:209-12
13. Ihara K, Toyada K, Ofuji A, Kawai S, Tuberculous bursitis of the greater trochanter. *J Orthop Sci*, 1998;3:120-4.

Conflict of Interest: Nil
Source of Support: None

How to Cite this Article:

Pimprikar MV, Kekatpure AL. Subdeltoid Bursa Tuberculosis with Rice bodies formation :Case Report and Review of literature. *Journal of Orthopaedic Case Reports* 2014 April-June;4(2): 57-59